


The modeling of the continuous emission spectrum of a dense non-ideal plasma in optical region


A. A. Mihajlov, N. M. Sakan, V. A. Sreković


VI Serbian Conference on Speactral Line Shapes in
Astrophysics 2007

The optical properties of dense plasma

The optical properties of dense plasma







$$U(r) = \begin{cases} -\frac{e^2}{r} + \frac{e^2}{r_c} & : & 0 < r \leq r_c \\ 0 & : & r_c < r \end{cases} \quad (1)$$

$$\left\{ \frac{d^2}{dr^2} + \frac{2}{r} \frac{d}{dr} + \frac{2m}{\hbar^2} [E - U(r)] - \frac{l(l+1)}{r^2} \right\} R = 0, \quad (2)$$

$$\left\{ \frac{d^2}{dr^2} + \frac{2}{r} \frac{d}{dr} + \frac{2m}{\hbar^2} [E - U(r)] - \frac{l(l+1)}{r^2} \right\} R = 0, \quad (2)$$

$$R(r) = \frac{P(r)}{r}, \quad (3)$$

$$\left\{ \frac{d^2}{dr^2} + \frac{2}{r} \frac{d}{dr} + \frac{2m}{\hbar^2} [E - U(r)] - \frac{l(l+1)}{r^2} \right\} R = 0, \quad (2)$$

$$R(r) = \frac{P(r)}{r}, \quad (3)$$

$$\frac{d^2 P(r)}{dr^2} + \left[\frac{2m}{\hbar^2} (E - U(r)) - \frac{l(l+1)}{r^2} \right] P(r) = 0, \quad (4)$$

$$\left\{ \frac{d^2}{dr^2} + \frac{2}{r} \frac{d}{dr} + \frac{2m}{\hbar^2} [E - U(r)] - \frac{l(l+1)}{r^2} \right\} R = 0, \quad (2)$$

$$R(r) = \frac{P(r)}{r}, \quad (3)$$

$$\frac{d^2 P(r)}{dr^2} + \left[\frac{2m}{\hbar^2} (E - U(r)) - \frac{l(l+1)}{r^2} \right] P(r) = 0, \quad (4)$$

The areas of the solution

$$\left\{ \frac{d^2}{dr^2} + \frac{2}{r} \frac{d}{dr} + \frac{2m}{\hbar^2} [E - U(r)] - \frac{l(l+1)}{r^2} \right\} R = 0, \quad (2)$$

$$R(r) = \frac{P(r)}{r}, \quad (3)$$

$$\frac{d^2 P(r)}{dr^2} + \left[\frac{2m}{\hbar^2} (E - U(r)) - \frac{l(l+1)}{r^2} \right] P(r) = 0, \quad (4)$$

The areas of the solution

a) $E < 0$

$$\left\{ \frac{d^2}{dr^2} + \frac{2}{r} \frac{d}{dr} + \frac{2m}{\hbar^2} [E - U(r)] - \frac{l(l+1)}{r^2} \right\} R = 0, \quad (2)$$

$$R(r) = \frac{P(r)}{r}, \quad (3)$$

$$\frac{d^2 P(r)}{dr^2} + \left[\frac{2m}{\hbar^2} (E - U(r)) - \frac{l(l+1)}{r^2} \right] P(r) = 0, \quad (4)$$

The areas of the solution

a) $E < 0$

b) $0 < E < e^2/r_c$

$$\left\{ \frac{d^2}{dr^2} + \frac{2}{r} \frac{d}{dr} + \frac{2m}{\hbar^2} [E - U(r)] - \frac{l(l+1)}{r^2} \right\} R = 0, \quad (2)$$

$$R(r) = \frac{P(r)}{r}, \quad (3)$$

$$\frac{d^2 P(r)}{dr^2} + \left[\frac{2m}{\hbar^2} (E - U(r)) - \frac{l(l+1)}{r^2} \right] P(r) = 0, \quad (4)$$

The areas of the solution

- a) $E < 0$
- b) $0 < E < e^2/r_c$
- c) $e^2/r_c < E$

The bond states, case a)

$$P(r) = \begin{cases} C_N M_{\kappa, \mu} \left(\frac{2r}{a_0 \nu^2} \right) & : r < r_c \\ \sqrt{r} C_N C_{E, l} k_{l+1/2} \left(r \sqrt{-2mE/\hbar^2} \right) & : r \geq r_c \end{cases} \quad (5)$$

Where $\mu = l + 1/2$, $\nu = \sqrt{-\frac{e^2}{a_0} \frac{1}{2E_c}}$, $a_0 = \hbar^2/me^2$ and $E_c = E + e^2/r_c < 0$.

The bond states, case a)

$$P(r) = \begin{cases} C_N M_{\kappa, \mu} \left(\frac{2r}{a_0 \nu^2} \right) & : r < r_c \\ \sqrt{r} C_N C_{E, l} k_{l+1/2} \left(r \sqrt{-2mE/\hbar^2} \right) & : r \geq r_c \end{cases} \quad (5)$$

Where $\mu = l + 1/2$, $\nu = \sqrt{-\frac{e^2}{a_0} \frac{1}{2E_c}}$, $a_0 = \hbar^2 / me^2$ and $E_c = E + e^2 / r_c < 0$.


Free states, case b)


$$P(r) = \begin{cases} C_0 M_{\kappa, \mu} \left(\frac{2r}{a_0 \nu^2} \right) & : r < r_c \\ C_0 r \left[C_1 \frac{\sqrt{2mE}}{\hbar} j_l \left(\frac{\sqrt{2mE}}{\hbar} r \right) + C_2 \frac{\sqrt{2mE}}{\hbar} y_l \left(\frac{\sqrt{2mE}}{\hbar} r \right) \right] & : r \geq r_c \end{cases} \quad (6)$$

Free states, case c)

$$P(r) = \begin{cases} C_0 F_l(\eta, r) & : r < r_c \\ C_0 r \left[C_1 \frac{\sqrt{2mE}}{\hbar} j_l \left(\frac{\sqrt{2mE}}{\hbar} r \right) + C_2 \frac{\sqrt{2mE}}{\hbar} y_l \left(\frac{\sqrt{2mE}}{\hbar} r \right) \right] & : r \geq r_c \end{cases} \quad (7)$$


Where $\eta = 1/\sqrt{2(E - 1/r_c)}$








Lorentz:


$$\frac{\gamma}{2\pi} \frac{1}{(E - E_n)^2 + \gamma^2/2} \quad (8)$$













$$U(r) = \begin{cases} -\frac{e^2}{r} + k_{rc} \frac{e^2}{r_c} & : \quad 0 < r \leq r_c \\ 0 & : \quad r_c < r \end{cases} \quad (9)$$



$$U(r) = \begin{cases} -\frac{e^2}{r} + k_{rc} \frac{e^2}{r_c} & : & 0 < r \leq r_c \\ 0 & : & r_c < r \end{cases} \quad (9)$$


The areas of the solution



$$U(r) = \begin{cases} -\frac{e^2}{r} + k_{rc} \frac{e^2}{r_c} & : & 0 < r \leq r_c \\ 0 & : & r_c < r \end{cases} \quad (9)$$

The areas of the solution

a) $E < 0$




$$U(r) = \begin{cases} -\frac{e^2}{r} + k_{rc}\frac{e^2}{r_c} & : & 0 < r \leq r_c \\ 0 & : & r_c < r \end{cases} \quad (9)$$

The areas of the solution

a) $E < 0$

b) $0 < E < k_{rc}e^2/r_c$



$$U(r) = \begin{cases} -\frac{e^2}{r} + k_{rc}\frac{e^2}{r_c} & : & 0 < r \leq r_c \\ 0 & : & r_c < r \end{cases} \quad (9)$$

The areas of the solution

- a) $E < 0$
- b) $0 < E < k_{rc}e^2/r_c$
- c) $k_{rc}e^2/r_c < E$

The bond states, case a)

$$P(r) = \begin{cases} C_N M_{\kappa, \mu} \left(\frac{2r}{a_0 \nu^2} \right) & : \quad r < r_c \\ \sqrt{r} C_N C_{E, l} k_{l+1/2} \left(r \sqrt{-2mE/\hbar^2} \right) & : \quad r \geq r_c \end{cases} \quad (10)$$

Where $\mu = l + 1/2$, $\nu = \sqrt{-\frac{e^2}{a_0} \frac{1}{2E_c}}$, $a_0 = \hbar^2/m e^2$ and $E_c = E + k_{rc} e^2/r_c < 0$.

The bond states, case a)

$$P(r) = \begin{cases} C_N M_{\kappa, \mu} \left(\frac{2r}{a_0 \nu^2} \right) & : r < r_c \\ \sqrt{r} C_N C_{E, l} k_{l+1/2} \left(r \sqrt{-2mE/\hbar^2} \right) & : r \geq r_c \end{cases} \quad (10)$$

Where $\mu = l + 1/2$, $\nu = \sqrt{-\frac{e^2}{a_0} \frac{1}{2E_c}}$, $a_0 = \hbar^2/me^2$ and $E_c = E + k_{rc}e^2/r_c < 0$.

Free states, case b)

$$P(r) = \begin{cases} C_0 M_{\kappa, \mu} \left(\frac{2r}{a_0 \nu^2} \right) & : r < r_c \\ C_0 r \left[C_1 \frac{\sqrt{2mE}}{\hbar} j_l \left(\frac{\sqrt{2mE}}{\hbar} r \right) + C_2 \frac{\sqrt{2mE}}{\hbar} y_l \left(\frac{\sqrt{2mE}}{\hbar} r \right) \right] & : r \geq r_c \end{cases} \quad (11)$$

Free states, case c)

$$P(r) = \begin{cases} C_0 F_l(\eta, r) & : r < r_c \\ C_0 r \left[C_1 \frac{\sqrt{2mE}}{\hbar} j_l \left(\frac{\sqrt{2mE}}{\hbar} r \right) + C_2 \frac{\sqrt{2mE}}{\hbar} y_l \left(\frac{\sqrt{2mE}}{\hbar} r \right) \right] & : r \geq r_c \end{cases} \quad (12)$$

Where $\eta = 1/\sqrt{2(E - k_{rc}1/r_c)}$

Example of a shift

The parameter k_{rc} is set to be $5/2$

n	l	E_1 [eV]	E_2 [eV]
1	0	-13.1107	-12.3687
2	0	-2.90672	-2.16467
2	1	-2.90645	-2.16467
3	0	-1.01716	-0.275107
3	1	-1.01716	-0.275107
3	2	-1.01683	-0.275047

Conclusions

Conclusions

- The cut-off Coulomb potential is a unique QM model that enables good agreement with experimental data in areas of the plasmas with mid and high nonideality.

Conclusions

- The cut-off Coulomb potential is a unique QM model that enables good agreement with experimental data in areas of the plasmas with mid and high nonideality.
- The possibility of connecting the optical and transport coefficients of plasma in the same QM figure.

Conclusions

- The cut-off Coulomb potential is a unique QM model that enables good agreement with experimental data in areas of the plasmas with mid and high nonideality.
- The possibility of connecting the optical and transport coefficients of plasma in the same QM figure.
- Flexible, easily includes additional processes in the QM figure.

Conclusions

- The cut-off Coulomb potential is a unique QM model that enables good agreement with experimental data in areas of the plasmas with mid and high nonideality.
- The possibility of connecting the optical and transport coefficients of plasma in the same QM figure.
- Flexibly, easily includes additional processes in the QM figure.
- Into the newly introduced QM model potential, the additional shifting of levels is a part of the model itself.