

On the Stark Broadening in Hot Stars

Z. Simić¹, M. S. Dimitrijević¹ & A. Kovačević²

1. Astronomical Observatory Belgrade, Volgina 7, 11160 Belgrade, Serbia

2. Department for Astronomy, Faculty for Mathematics, University of Belgrade, Studentski Trg 16, 11000 Belgrade, Serbia

e-mail: zsimic@aob.bg.ac.rs, mdimitrijevic@aob.bg.ac.yu, andjelka@matf.bg.ac.rs

Abstract

In hot star atmospheres exist conditions where Stark widths are comparable and even larger than the thermal Doppler widths, so that the corresponding line broadening parameters are of importance for the hot star plasma investigation. Here, we investigated theoretically the influence of collisions with charged particles on heavy metal spectral line profiles for Te I, Cr II and Sn III in spectra of A stars and white dwarfs. We applied semiclassical perturbation theory. When it can not be applied in an adequate way, due to the lack of reliable atomic data, we used modified semiempirical theory.

We presented Stark broadening parameters, widths and shifts, for four Te I spectral lines, four resonant Cr II $3d^5-3d^4p$ spectral lines and two Sn III spectral lines. In the case with the available experimental and other theoretical data for the considered spectral lines we analyzed the agreement or a disagreement with our theoretical results. Also, here we considered the contributions of different collisional processes to the total Stark width in comparison with Doppler one.

We made a detailed analyzes of obtained results in order to compare Doppler and Stark broadening contributions in hot star atmospheres. We found that Stark broadening mechanism may be important in such plasma conditions and should be taken into account.

On the Stark Broadening of Cu I Spectral Lines

B. Zmerli¹, N. Ben Nessib¹, M. S. Dimitrijević^{2,3} & S. Sahal-Bréchot³

¹Groupe de Recherche en Physique Atomique et Astrophysique, INSAT, Centre Urbain Nord B. P. No. 676, 1080 Tunis, Cedex, Tunisia

²Astronomical Observatory, Volgina 7, 11060 Belgrade 38, Serbia

³Observatoire de Paris, 92195 Meudon, Cedex, France

e-mail: nebil.benessib@planet.tn, mdimitrijevic@aob.bg.ac.yu, sylvie.sahal-brechot@obspm.fr

Abstract

We investigate here the Stark broadening parameters of neutral copper spectral lines. This metal is often used in electrical industry as electrode materials, so that the data on its spectral lines are important not only for plasma research but also for diagnostic techniques in industrial laboratories.

Recently, temperature dependence of Stark width for neutral atom spectral lines is investigated (Zmerli et al., 2008), in order to find a method for scaling of Stark broadening parameters with temperature, better than the dependence $T^{1/2}$. In this work, Stark width dependence on T is analyzed using the lines of neutral helium.

Here, we calculated, using semiclassical perturbation theory of Sahal-Bréchot Stark widths and shifts for CuI 324.75, 327.39, 510.54, 515.32, 521.82 and

578.21 nm spectral lines. Our results are compared with different available experimental and theoretical data. Also, they are used to study the dependence of Stark broadening parameters with temperature.

References

Zmerli, B., Ben Nessib, N., Dimitrijević, M. S.: 2008, *European J. Phys. D*, **48**, 389.