

ANALYSIS OF STARK BROADENING PARAMETERS OF N II SPECTRAL LINES

M. Christova¹, M. S. Dimitrijević^{2,3}  and S. Sahal-Bréchet³

¹*Department of Applied Physics, Technical University of Sofia,
1000 Sofia, Bulgaria*

²*Astronomical Observatory, Volgina 7, 11060 Belgrade, Serbia*

³*LERMA, Observatoire de Paris, Université PSL, CNRS, Sorbonne Université,
F-92190 Meudon, France*

E-mail: mchristo@tu-sofia.bg

mdimitrijevic@aob.rs

sylvie.sahal-brechet@obspm.fr

We propose a study of Stark broadening parameters of N II spectral lines. The conditions of interest cover wide temperature interval (from 5 000 K to 200 000 K) and a perturber density of 10^{17} cm⁻³, suitable for astrophysical applications, laboratory diagnostics and for proton-boron fusion plasma research. The interactions that broad the spectral lines include particles as electrons, protons, alpha particles and boron ions in different stages of ionization. Recently, many experimental and theoretical studies are dedicated to proton-boron nuclear fusion which delivers energy with several significant advantages (Belloni, 2022).

References

Belloni, F.: 2022, Multiplication Processes in High-Density H-11B Fusion Fuel., Laser and Particle Beams, 3952779.

Acknowledgments: This work has been supported with a STSM visit grant for M.S.D. within the framework of COST Action CA21128-PROBONO “Proton Boron Nuclear Fusion: From Energy Production To Medical Applications”, supported by COST (European Cooperation in Science and Technology-www.cost.eu).