

PROBING BLR KINEMATICS VIA DOUBLE-PEAKED CA II

M. W. Ochmann¹, P. M. Weilbacher², M. A. Probst¹,
W. Kollatschny¹, D. Chelouche^{3, 4}, R. Chini⁵, D. Grupe⁶,
M. Haas⁵, S. Kaspi⁷ and S. Komossa⁸

¹*Institut für Astrophysik und Geophysik, Universität Göttingen,
Friedrich-Hund Platz 1, 37077 Göttingen, Germany*

²*Leibniz-Institut für Astrophysik Potsdam (AIP),
An der Sternwarte 16, 14482 Potsdam, Germany*

³*Department of Physics, Faculty of Natural Sciences, University of Haifa
Haifa 3498838, Israel*

⁴*Haifa Research Center for Theoretical Physics and Astrophysics, University of Haifa,
Haifa 3498838, Israel*

⁵*Ruhr University Bochum, Faculty of Physics and Astronomy,
Astronomical Institute (AIRUB), 44780 Bochum, Germany*

⁶*Department of Physics, Geology, and Engineering Technology, Northern
Kentucky University, 1 Nunn Drive, Highland Heights, KY 41099, USA*

⁷*School of Physics & Astronomy and the Wise Observatory,
The Raymond and Beverly Sackler Faculty of Exact Sciences,
Tel-Aviv 6997801, Israel*

⁸*Max-Planck-Institut für Radioastronomie,
Auf dem Hügel 69, D-53121 Bonn, Germany*

E-mail: martin.ochmann@uni-goettingen.de

Double-peaked emission lines are observed in a small fraction of active galactic nuclei (AGN) and offer a means to determine key properties of the line-emitting region, known as the broad-line region (BLR). Until now, studies of double-peaked emitters (DPEs) have focused almost exclusively on the Balmer lines H β and H α , while investigations of double-peaked lines in the UV or near-infrared remain scarce. Notably, we recently reported – for the first time – broad double-peaked emission profiles in the near-infrared Ca II triplet for individual sources, namely NGC 1566 and NGC 4593. These

detections represent the first of their kind to date. The profiles enabled us to link the kinematics of the Ca II-emitting phase of the BLR to those of a rotating relativistic disk, largely unaffected by internal turbulence, thus allowing a more direct interpretation of the underlying BLR dynamics. In this talk, we summarize our recent findings and discuss the potential of Ca II emission profiles as a powerful diagnostic tool for probing the kinematics of the BLR.