

REFINING BLACK HOLE MASS ESTIMATES IN TYPE 1 AGN: DEPENDENCE ON $H\beta$ LINE PROPERTIES

S. Marčeta-Mandić¹ , J. Kovačević-Dojčinović¹  and
L. Č. Popović^{1,2,3} 

¹*Astronomical Observatory, Volgina 7, 11060 Belgrade, Serbia*

²*Faculty of Mathematics - University of Belgrade, Department of Astronomy,
Studentski trg 16, 11000 Belgrade, Serbia*

³*PIFI Research Fellow, Key Laboratory for Particle Astrophysics,
Institute of High Energy Physics, Chinese Academy of Sciences,
19B Yuquan Road, 100049 Beijing, China*

E-mail: sladjana@aob.rs

Active Galactic Nuclei (AGN) are compact regions at the centers of active galaxies that exhibit a pronounced excess in luminosity, typically attributed to the accretion of gas onto a central supermassive black hole (SMBH). Reliable estimates of SMBH masses are essential for understanding both AGN physics and galaxy evolution. In this study, we explore the robustness of black hole mass estimates derived from the broad $H\beta$ emission line in Type 1 AGN spectra, using a sample from the Sloan Digital Sky Survey (SDSS). We compare these virial mass estimates with stellar velocity dispersion (σ_*) as an independent tracer of supermassive black hole mass. Our analysis reveals that the correlation between $H\beta$ -based masses and σ_* significantly improves in specific spectral sub-samples, namely those with broader $H\beta$ profiles, red-ward asymmetry, high continuum luminosity, and no evident signatures of outflows. These results indicate that, in AGN exhibiting such spectral characteristics, the $H\beta$ line can serve as a more reliable virial mass estimator. This has important implications for the selection of AGN sub-populations in future spectroscopic surveys and for refining empirical scaling relations used in SMBH mass determinations.