## Rate coefficients of He<sup>+</sup> ions in CF<sub>4</sub> gas

## Željka D. Nikitović and Zoran M. Raspopović

Institute of Physics, University of Belgrade, Pregrevica 118, 11080 Belgrade, Serbia E-mail: zeljka@ipb.ac.rs

This paper is dedicated to the presentation of the set of  $He^+$  ions scattering cross-sections in  $CF_4$  which is estimated using available experimental data for exothermic charge transfer cross-sections producing  $CF_3^+$  and  $CF_2^+$  ions and endothermic charge transfer cross-section producing  $CF^+$ ,  $C^+$  and  $F^+$  ions. Due to significant particle losses, the experimental transport coefficients were not measured. The transport properties of  $He^+$  ions in  $CF_4$  required to model the discharge containing the mentioned ions were calculated by the Monte Carlo method at a temperature of T=300 K. In this paper we give the characteristic energy and specially rate coefficients for low and moderately reduced electric fields E /N (E-electric field, N-gas density) and accounting for non-conservative collisions.

He-CF<sub>4</sub> mixtures are used in gas electron multipliers for various imaging purposes (X-rays, charged particles, thermal neutrons and dark matter detection) (Fraga M.M.F.R. et al. 2008). Bursts of electron multiplication affect production of various ions that may affect time distribution of detected particles (Bošnjaković D. 2016).

## References

Fraga M.M.F.R., Fraga F.A.F., Fetal S.T.G., Margato L.M.S., Ferreira Marques R., Policarpo A.J.P.L., 2003, Nucl. Instrum. Meth. in Phys. Res. A 504, 88; Kaboth A., Monroe J., Ahlen S., Dujmić D., Henderson S., Kohse G., Lanza R., Lewandowska M., Roccaro A., Sciolla G., Skvorodnev N., Tomita H., Vanderspek R., Wellenstein H., Yamamoto R., Fisher P., 2008, Nucl. Instrum. and Meth.in Phys.Res. A 592, 63.

Bošnjaković D.,2016, Ph Dissertation, Faculty of Electrical Engineering, University of Belgrade, Belgrade, Serbia.