

BH MASS ESTIMATION for Quasars

J. Sulentic -- IAA - CSIC
P. Marziani -- OA Padova - INAF

Estimating BH Mass

Broad Emission Lines as virial estimators

FWHM Hbeta → virial v

Single epoch FWHM measures? rms? Composites?
Or sigma?

$$M_{\rm BH} = \frac{v^2 r_{\rm BLR}}{G}$$

$$v=f{\rm FWHM}$$

$$v=\sqrt{3}/2~{\rm FWIIM}(\Pi\beta_{\rm BC})$$

The f factor?


- BLR geometry and kinematics
- Strong inclination dependence
- Of order $\sim\sim 1-2$ (or 5.5 Onken et al. 2009 using sigma)
- Same for Pop. A and B?

Not all FWHM Hbeta are the same


Sources above and below FWHM
Hbeta=4000km/s show different
structure (Sulentic et al. 2002)

Relationship between FWHM and sigma changes --
reflecting multiple components
(Collin et al. 2006: $f(A) \approx 2.12$ $f(B) \approx 0.5-1.0$)

Average Quasar Spectrum: Francis et al. 1991




“Eigenvectors of Quasars”




✿ Quasar spectra are
not similar!

↳ historic inability to explain
line ratios via simple
photoionization models



Sulentic et al. 2002:
LIL/BLR Structural
Difference between
Population A and B
(Sample of about 200
Seyfert 1 and
low-redshift quasars
High S/N and resolution
4 Å FWHM)






Extreme M BH


- Pop. A extremely narrow profiles lead to underestimates of MBH
- Most extreme MBH values came from pop B sources using FWHM Hbeta uncorrected for the extra very broad component.

Above z \sim 0.7?

- Follow H β into the IR to z = 3.8
- MgII2798 with suitable corrections to z > 6
- CIV1549 dangerous







So we have “v” – ways to estimate BLR size


- Directly: reverberation mapping
(Peterson & Horne 2006)
- Using r_BLR vs. Luminosity relation
(Marziani et al. 2009; Trachtenbrot et al. 2011)
- Photoionization Method
(Dibai 1984; Wandel & Yahil 1985; Padovani & Rafanelli 1989;
Wandel et al. 1999; Negrete et al. 2011)

Determination of central compact object mass

$c\tau_{\max}$ provides an emissivity weighted estimate of the BLR linear distance (size) from central continuum source

$$r_{BLR} \approx c\tau_{\max} \approx 33 \left(\frac{\lambda L_{5100}}{10^{44} \text{ erg s}^{-1}} \right)^{0.7} \text{ l.d.}$$

Kaspi et al. 2000



More recent work indicates exponent ≈ 0.5

Bentz et al. 2009


Paves the road to easy mass estimates

Bolometric luminosity estimated from bolometric correction $\approx 9 - 10$ times the optical luminosity (at 5100 Å)

Elvis et al. 1994; see Nemmen & Brotherton 2010
for a more modern approach

$$U = \frac{\int_{\nu_0}^{+\infty} \frac{L_\nu}{h\nu} d\nu}{4\pi n_{\text{H}} c r^2}$$

$$r_{\text{BLR}} = \underbrace{\frac{1}{(4\pi c)^{\frac{1}{2}}}}_{\text{const.}} \underbrace{(Un_{\text{H}})^{-\frac{1}{2}}}_{\text{diagnostics}} \left(\underbrace{\int_{\nu_0}^{+\infty} \frac{L_\nu}{h\nu} d\nu}_{\# \text{ ionizing photons}} \right)^{\frac{1}{2}}$$




$$M_{\rm BH} = \frac{3}{4}\frac{r_{\rm BLR}{\rm FWHM(H}\beta_{\rm BC})^2}{G}$$

$$r_{\rm BLR} \propto (L_{5100})^\alpha \qquad (Kaspi \; et \; al., \; 2000; \; 2005)$$

$$r_{BLR} = 0.85 \cdot 10^{17} \cdot \left[\frac{\lambda L_{\lambda}\left(5100\text{\AA}\right)}{10^{44}\,\mathrm{erg\,s^{-1}}} \right]^{0.7} \mathrm{cm}$$

$$\lambda L_{\lambda}\left(5100\text{\AA}\right) = 3.14 \cdot 10^{35 - 0.4(M_B)}$$

$$\lambda L_{\lambda}\left(5100\text{\AA}\right) = 4\pi d_{\rm P}^2 \lambda f_{\lambda} {\rm ergs~s^{-1}}$$






Table 6. Measurements on the broad lines of median spectra.

Object name (1)	$W(H\beta)^a$ (2)	$W(FeII\lambda 4570)^a$ (3)	$FWHM(FeII)^b$ (4)	$F(H\beta_{BC})/F(H\beta)^c$ (5)	$FWHM(H\beta_{BC})^d$ (6)	$\log M_{BH} H\beta_{BC}^e$ (7)	$\log L_{bol}/L_{Edd} H\beta_{BC}^e$ (8)
A1	72 ⁺¹¹ ₋₁₁	26 ⁺³ ₋₄	2700 ⁺¹¹⁰⁰ ₋₁₁₀₀	1.00
A2	65 ⁺¹⁰ ₋₁₂	49 ⁺¹³ ₋₁₁	3700 ⁺²⁰⁰⁰ ₋₁₄₀₀	1.00
B1	86 ⁺¹³ ₋₁₃	26 ⁺⁵ ₋₆	5200 ⁺²⁴⁰⁰ ₋₂₃₀₀	0.27	4000
B2	70 ⁺¹¹ ₋₁₁	44 ⁺⁸ ₋₁₄	5000 ⁺⁸⁰⁰ ₋₁₇₀₀	0.32	4000
A	61 ⁺¹⁰ ₋₁₃	25 ⁺⁷ ₋₇	2700 ⁺¹⁵⁰⁰ ₋₁₂₀₀	1.00
M	67 ⁺¹⁰ ₋₁₁	35 ⁺⁷ ₋₆	3800 ⁺¹⁴⁵⁰ ₋₁₂₀₀	1.00
MB	86 ⁺¹⁰ ₋₁₃	31 ⁺⁶ ₋₇	5000 ⁺¹⁶⁰⁰ ₋₁₈₀₀	0.27	4100
43A	91 ⁺¹⁰ ₋₂₀	36 ⁺⁷ ₋₇	3000 ⁺⁵⁰⁰ ₋₇₅₀	1.00	...	6.1	-0.74
44A	69 ⁺¹³ ₋₁₅	38 ⁺¹⁰ ₋₁₀	2600 ⁺⁵⁰⁰ ₋₇₅₀	1.00	...	6.8	-0.47
45A	86 ⁺¹⁰ ₋₂₀	43 ⁺¹⁰ ₋₁₀	2800 ⁺⁷⁵⁰ ₋₅₀₀	1.00	...	7.8	-0.43
46A	80 ⁺¹⁰ ₋₁₉	47 ⁺¹⁰ ₋₁₀	3000 ⁺⁶⁰⁰ ₋₆₀₀	1.00	...	8.6	-0.26
47A	68 ⁺¹⁰ ₋₁₁	30 ⁺⁸ ₋₈	3000 ⁺¹⁴⁰⁰ ₋₁₂₀₀	1.00	...	9.6	-0.20
48A	60 ⁺¹¹ ₋₁₁	27 ⁺⁵ ₋₈	3800 ⁺¹⁵⁰⁰ ₋₁₁₀₀	1.00	...	10.3	+0.11
43B	130 ⁺²⁰ ₋₂₀	8 ⁺¹⁰ ₋₇	...	0.59	4600	7.1	-0.68
44B	125 ⁺¹⁸ ₋₃₉	38 ⁺⁵ ₋₂₀	5600 ⁺⁶⁰⁰ ₋₁₈₀₀	0.49	4700	7.7	-1.37
45B	111 ⁺¹³ ₋₂₀	29 ⁺⁵ ₋₁₅	4900 ⁺⁵⁰⁰ ₋₈₀₀	0.35	4400	8.4	-0.98
46B	93 ⁺¹⁰ ₋₂₀	22 ⁺⁵ ₋₁₀	5900 ⁺³⁵⁰ ₋₁₂₀₀	0.37	4800	9.1	-0.73
47B	92 ⁺¹³ ₋₁₄	38 ⁺⁷ ₋₇	4900 ⁺¹⁶⁰⁰ ₋₂₀₀₀	0.27	4000	9.6	-0.24
48B	75 ⁺⁹ ₋₁₁	12 ⁺³ ₋₃	4600 ⁺¹²⁰⁰ ₋₁₇₀₀	0.23	4300	10.3	+0.03

^a Equivalent width of H β ($H\beta_{BC} + H\beta_{VBC}$) and FeII $\lambda 4570$ in Å $\pm 2\sigma$ confidence level uncertainty. Note that those values have been computed on median spectra with flux normalized to unity at $\lambda = 5100$ Å. Considering that the continuum shape is not flat, but that there is however little dispersion in continuum shape across the median spectra, it is $W(H\beta) \approx I(H\beta)/1.1$, and $W(FeII\lambda 4570) \approx I(FeII\lambda 4570)/1.25$ Å. ^b FWHM of lines in the blend in units of km s⁻¹ computed by specfit as for the individual sources. Uncertainty is at $\pm 2\sigma$ confidence level. See text for details. ^c Intensity ratio of the H β_{BC} to total H β line emission i.e., H β_{BC} and H β_{VBC} . ^d FWHM of the H β_{BC} component i.e., after removing H β_{VBC} .

^e Logarithm of M_{BH} , in solar masses, and of L_{bol}/L_{Edd} . Values have been computed following Paper II, using the $FWHM(H\beta_{BC})$ reported in Col. (6), and assuming the average bin luminosity. Values are therefore only indicative. No M_{BH} or L_{bol}/L_{Edd} has been computed for median in spectral types since they are normalized median spectra made regardless of their luminosity. ^f FeII $_{opt}$ too faint for FWHM to be meaningfully constrained.

