

On the correlation between line-of-sight velocity and metallicity for nearby F, G, K stars

Sonja Vidojević & Slobodan Ninković


sonja@matf.bg.ac.yu sninkovic@aob.bg.ac.yu

**VI Serbian Conference on Spectral Line Shapes in Astrophysics
Sremski Karlovci, Serbia, June 11-15 2007**

Treated data

- Spectroscopic data from SPOCS
(<http://vizier.u-strasbg.fr/viz-bin/VizieR>)
- Valenti J. A., Fischer D. A., 2005,
ApJS, 159, 141
- 1040 stars including the Sun

Line-of-sight velocity versus metallicity for all 1039 SPOCS stars.



Percentages of galactic subsystems

- thin disc 93%
- thick disc 6%
- halo 1 %


Thin disc

- ☐ Velocity ellipsoid determined on the bases of line-of-sight velocities

v_\odot	L_\odot	B_\odot	σ_U	σ_V	σ_W	l_V
25	70°	19°	34	22	17	6°

- First three yield Solar motion and the last one is vertex deviation

Age-metallicity relation



Thick disc

- On the average less metallic than thin disc
- Higher heliocentric velocities than in thin disc
- On the average age seems to be higher than in thin disc

Halo

- Separated most clearly
- Halo stars in this sample least metallic ≤ -0.8
- Highest moduli of line-of-sight velocities

References

- Alcobé S., Cubarsi R., 2005, A&A, 442, 929
- Dehnen W., Binney J. J., 1998, MNRAS, 298, 387
- Famaey B., Jorissen A., Luri X., Mayron M., Udry S., Dejonghe H., Turon C., 2005, A&A, 430, 165
- Nissen P. E., Schuster W. J., 1991, A&A, 251, 457
- Nordström B., Mayor M., Andersen J., Holmberg J., Pont F., Jørgensen B. R., Olsen E. H., Udry S., Mowlavi N., 2004, A&A, 418, 989
- Soubiran C., Girard P., 2005, A&A, 438, 139
- Valenti J. A., Fischer D. A., 2005, ApJS, 159, 141